Author Index

Abe, H., Tamura, S. and Kondo, H., ocalization of mRNA for protein phosphatase 2C in the brain of adult rats, 283 Agarwal, B.L., see La Gamma, E.F., 189 Altar, C.A., see Iacopino, A., 251 Andreussi, L., see Stenger, A.M., 273 Angelides, K.J., see Wang, C., 289

Banerjee, S., see Wong, M.-L., 19
Banker, G., see Kleiman, R., 103
Baraban, J.M., see Bhat, R.V., 263
Barraclough, C.A., see Hartman,
R.D., 223

R.D., 223
Barraclough, C.A., see Liaw, J.-J., 171
Bendotti, C., see Kruger, L., 267
Bergeron, C., see Sutherland, M.K., 239
Bertherat, J., see Llorens-cortes, C., 277
Bhat, A.N., see Kindy, M.S., 199
Bhat, N.R., see Kindy, M.S., 199

Bhat, R.V., Worley, P.F., Cole, A.J. and Baraban, J.M., Activation of the zinc finger encoding gene *krox*-20 in adult rat brain: comparison with *zif*268, 263

Bilkey, D.K., see Dragunow, M., 119Bixby, J.L., Identification of an alternatively spliced avian member of the synaptophysin gene family, 339

Brisigotti, M., see Stenger, A.M., 273 Brown, E.E., see Hurd, Y.L., 165 Brown, I.R., see Ni, B., 7 Bruce, J., see Shneidman, P.S., 127 Bruinvels, A.T., see Palacios, J.M., 175

Callea, F., see Stenger, A.M., 273
 Cama, A., see Stenger, A.M., 273
 Carpenter, M.K., Parker, I. and Miledi, R., Messenger RNAs coding for receptors and channels in the cerebral cortex of adult and aged rats, 1

Christakos, S., see Iacopino, A., 251
Cole, A.J., see Bhat, R.V., 263
Collinge, J., see Owen, F., 155
Crapper McLachlan, D.R., see Sutherland, M.K., 239

Crow, T.J., see Owen, F., 155

De Haan, J.B., Newman, J.D. and Kola, I., Cu/Zn superoxide dismutase mRNA and enzyme activity, and susceptibility to lipid peroxidation, increases with aging in murine brains, 179

DeCristofaro, J.D., see La Gamma, E.F., 189

DiStefano, P.S., see Matheny, C., 75 Doi, S.Q., see Wong, M.-L., 19 Donovan, D.M., see Takemura, M., 207

Dragunow, M., see Hughes, P., 355
Dragunow, M., Yamada, N., Bilkey, D.K.
and Lawlor, P., Induction of immediateearly gene proteins in dentate granule
cells and somatostatin interneurons after
hippocampal seizures, 119

Epelbaum, J., see Llorens-cortes, C., 277

Fabbretti, G., see Stenger, A.M., 273

Ferri, G.-L., Levi, A. and Possenti, R., A novel neuroendocrine gene product: selective VGF8a gene expression and immuno-localisation of the VGF protein in endocrine and neuronal populations, 139

Fibiger, H.C., see Hurd, Y.L., 165 Finch, C.E., see Pasinetti, G.M., 63 Finlay, J.M., see Hurd, Y.L., 165

Garré, M.L., see Stenger, A.M., 273 German, D., see Iacopino, A., 251 Gerfen, C.R., see Hurd, Y.L., 165 Gluckman, P.D., see Klempt, N.D., 93 Gold, P.W., see Wong, M.-L., 19 Gundlach, A.L., see Rutherfurd, S.D., 301 Gunn, A.J., see Klempt, N.D., 93 Gurd, J.W., see Ni, B., 7

Hammond, C., see Morton, A.J., 53
Harding, A.E., see Owen, F., 155
Hartman, R.D., Liaw, J.-J., He, J.-R. and
Barraclough, C.A., Effects of reserpine
on tyrosine hydroxylase mRNA levels in
locus coeruleus and medullary A1 and
A2 neurons analyzed by in situ hybridization histochemistry and quantitative
image analysis methods, 223

Hartman, R.D., see Liaw, J.-J., 231 Haussler, M.R., see Sutherland, M.K., 239 He, J.-R., see Hartman, R.D., 223 He, J.-R., see Liaw, J.-J., 171 Henderson, G., see Morton, A.J., 53 Hidaka, H., see Morita, S., 313

Hoyer, D., see Palacios, J.M., 175
Hughes, P., Lawlor, P. and Dragunow, M.,
Basal expression of Fos, Fos-related,
Jun, and Krox 24 proteins in rat hippocamous. 355

Hurd, Y.L., Brown, E.E., Finlay, J.M., Fibiger, H.C. and Gerfen, C.R., Temporal changes in tyrosine hydroxylase mRNA levels in A1, A2 and locus ceruleus neurons following electrical stimulation of A1 noradrenergic neurons, 165

Iacopino, A., Christakos, S., German, D., Sonsalla, P.K. and Altar, C.A., Calbindin-D_{28K}-containing neurons in animal models of neurodegeneration: possible protection from excitotoxicity. 251

Iismaa, T.P., see Selbie, L.A., 159 Ishii, D.N., see Wang, C., 289

Jomary, C., see Llorens-cortes, C., 277

Kawai, N., see Shimazaki, K., 331
Kelly, A.B., see Pasinetti, G.M., 63
Kindy, M.S., Bhat, A.N. and Bhat, N.R.,
Transient ischemia stimulates glial fibrillary acid protein and vimentin gene expression in the gerbil neocortex, striatum and hippocampus, 199
Kitayama, S., see Shimada, S., 359

Kleiman, R., Banker, G. and Steward, O., Temperature-dependent blockade of nucleocytoplasmic transport of newly synthesized RNA in neurons, 103

Klempt, M., see Klempt, N.D., 93
 Klempt, N.D., Sirimanne, E., Gunn, A.J.,
 Klempt, M., Singh, K., Williams, C. and
 Gluckman, P.D., Hypoxia-ischemia induces transforming growth factor β₁
 mRNA in the infant rat brain, 93

Kobayashi, K., see Morita, S., 313 Kohama, S., see Pasinetti, G.M., 63 Kola, I., see De Haan, J.B., 179 Kondo, H., see Abe, H., 283

Kordon, C., see Llorens-cortes, C., 277
Kruger, L., Bendotti, C., Rivolta, R. and
Samanin, R., GAP-43 mRNA localization in the rat hippocampus CA₃
field, 267

La Gamma, E.F., Agarwal, B.L. and De-Cristofaro, J.D., Regulation of adrenomedullary preproenkephalin mRNA: effects of hypoglycemia during development, 189

Lad, R., see Wong, M.-L., 19 Lawlor, P., see Dragunow, M., 119 Lawlor, P., see Hughes, P., 355 Leach, M., see Owen, F., 155 Levi, A., see Ferri, G.-L., 139 Li, Y., see Wang, C., 289

Liaw, J.-J., He, J.-R. and Barraclough, C.A., Temporal changes in tyrosine hydroxylase mRNA levels in A1, A2 and locus ceruleus neurons following electrical stimulation of A1 noradrenergic neurons. 171

Liaw, J.-J., see Hartman, R.D., 223 Licinio, J., see Wong, M.-L., 19

Lightman, S.L., see Young III, W.S., 111 Lindefors, N., Amphetamine and haloperidol modulate preprotachykinin A mRNA expression in rat nucleus accumbens and caudate-putamen, 151

Llorens-cortes, C., Bertherat, J., Jomary, C., Kordon, C. and Epelbaum, J., Regulation of somatostatin synthesis by GABA_A receptor stimulation in mouse brain, 277

Lofthouse, R., see Owen, F., 155 Louis, W.J., see Rutherfurd, S.D., 301

Martino, D.D., see Stenger, A.M., 273
 Mason, W.T., see Morton, A.J., 53
 Matheny, C., DiStefano, P.S. and Milbrandt, J., Differential activation of NGF receptor and early response genes in neural crest-derived cells, 75

McKinney, M., see Robbins, M., 83 Memo, M., see Valerio, A., 145 Mengod, G., see Rocamora, N., 27 Micevych, P.E., see Popper, P., 43 Milbrandt, J., see Matheny, C., 75 Miledi, R., see Campenter, M.K., 1

Miledi, R., see Carpenter, M.K., 1 Morgan, D.G., see Pasinetti, G.M., 63 Morita, S., Kobayashi, K., Hidaka, H. and Nagatsu, T., Organization and complete nucleotide sequence of the gene encoding mouse phenylethanolamine N-methyltransferase, 313

Morton, A.J., Hammond, C., Mason, W.T. and Henderson, G., Characterisation of the L- and N-type calcium channels in differentiated SH-SY5Y neuroblastoma cells: calcium imaging and single channel recording, 53

Nagatsu, T., see Morita, S., 313
Nakajima, T., see Shimazaki, K., 331
Newman, J.D., see De Haan, J.B., 179
Ni, B., Rush, S., Gurd, J.W. and Brown,
I.R., Molecular cloning of calmodulin
mRNA species which are preferentially
expressed in neurons in the rat brain, 7

Osterburg, H.H., see Pasinetti, G.M., 63 Owen, F., Poulter, M., Collinge, J., Leach, M., Lofthouse, R., Crow, T.J. and Harding, A.E., A dementing illness associated with a novel insertion in the prion protein gene, 155

Palacios, J.M., see Rocamora, N., 27
Palacios, J.M., Waeber, C., Bruinvels, A.T.
and Hoyer, D., Direct visualization of
serotonin_{1D} receptors in the human
brain using a new iodinated radioligand, 175

Parker, I., see Carpenter, M.K., 1
Pasinetti, G.M., Osterburg, H.H., Kelly,
A.B., Kohama, S., Morgan, D.G., Reinhard Jr., J.F., Stellwagen, R.H. and
Finch, C.E., Slow changes of tyrosine
hydroxylase gene expression in dopaminergic brain neurons after neurotoxin
lesioning: a model for neuron aging, 63

Pollack, A.E. and Wooten, G.F., D₂ dopaminergic regulation of striatal preproenkephalin mRNA levels is mediated at least in part through cholinergic interneurons, 35

Popper, P., Ulibarri, C. and Micevych, P.E., The role of target muscles in the expression of calcitonin gene-related peptide mRNA in the spinal nucleus of the bulbocavernosus, 43

Possenti, R., see Ferri, G.-L., 139 Post, R.M., see Wong, M.-L., 19 Poulter, M., see Owen, F., 155

Reinhard, Jr., J.F., see Pasinetti, G.M., 63 Rivolta, R., see Kruger, L., 267

Robbins, M. and McKinney, M., Transcriptional regulation of neuromodulin (GAP-43) in mouse neuroblastoma clone N1E-115 as evaluated by the RT-/PCR method, 83

Robinson, H.P.C., see Shimazaki, K., 331 Rocamora, N., Palacios, J.M. and Mengod, G., Limbic seizures induce a differential regulation of the expression of nerve growth factor, brain-derived neurotrophic factor and neurotrophin-3, in the rat hippocampus, 27

Rudeen, P.K., see Zoeller, R.T., 321 Rush, S., see Ni, B., 7

Rutherfurd, S.D., Widdop, R.E., Sannajust, F., Louis, W.J. and Gundlach, A.L., Expression of c-fos and NGFI-A messenger RNA in the medulla oblongata of the anaesthetized rat following stimulation of vagal and cardiovascular afferents, 301

Salvaterra, P.M., see Tajima, Y., 213
Samanin, R., see Kruger, L., 267
Sannajust, F., see Rutherfurd, S.D., 301
Schlaepfer, W.W., see Shneidman, P.S., 127
Schwartz, M.L., see Shneidman, P.S., 127
Selbie, L.A., Townsend-Nicholson, A., Iismaa, T.P. and Shine, J., Novel G
protein-coupled receptors: a gene family of putative human olfactory receptor sequences, 159

Shimada, S., Kitayama, S., Walther, D. and Uhl, G., Dopamine transporter mRNA: dense expression in ventral midbrain neurons, 359

Shimazaki, K., Robinson, H.P.C., Nakajima, T., Kawai, N. and Takenawa, T., Purification of AMPA type glutamate receptor by a spider toxin, 331

Shine, J., see Selbie, L.A., 159Shneidman, P.S., Bruce, J., Schwartz, M.L.and Schlaepfer, W.W.. Negative regulatory regions are present upstream in the

three mouse neurofilament genes, 127
Singh, K., see Klempt, N.D., 93
Sirimanne, E., see Klempt, N.D., 93
Smith, M.A., see Wong, M.-L., 19
Somerville, M.J., see Sutherland.

Somerville, M.J., see Sutherland M.K., 239 Sonsalla, P.K., see Iacopino, A., 251 Spano, P., see Valerio, A., 145

Stellwagen, R.H., see Pasinetti, G.M., 63
Stenger, A.M., Garré, M.L., Andreussi,
L., Cama, A., Callea, F., Brisigotti, M.,
Fabbretti, G., Martino, D.D. and Tonini,
G.P., Expression of histone H3 cell cycle-related gene, Vimentin and MYC genes in pediatric brain tumors. A preliminary analysis showing the different malignant cell growth potential, 273

Steward, O., see Kleiman, R., 103 Sumikawa, K., equences on the N-terminus of ACh receptor subunits regulate their assembly, 349

Sutherland, M.K., Somerville, M.J., Yoong, L.K.K., Bergeron, C., Haussler, M.R. and Crapper McLachlan, D.R., Reduction of vitamin D hormone receptor mRNA levels in Alzheimer as compared to Huntington hippocampus: correlation with calbindin-28k mRNA levels, 239

Tajima, Y. and Salvaterra, P.M., Positive and negative feedback regulation of choline acetyltransferase mRNA levels in *Drosophila*: a study using temperature-sensitive mutants and embryo cell cultures. 213

Takemura, M., Donovan, D.M. and Uhl, G.R., Primary afferent stimulation acts through a 193 base pair promoter region to upregulate preproenkephalin expression in dorsal horn of transgenic mice, 207

Takenawa, T., see Shimazaki, K., 331
Tamura, S., see Abe, H., 283
Tinti, C., see Valerio, A., 145
Tonini, G.P., see Stenger, A.M., 273
Townsend-Nicholson, A., see Selbie, L.A., 159

Uhl, G., see Shimada, S., 359 Uhl, G.R., see Takemura, M., 207 Ulibarri, C., see Popper, P., 43

Valerio, A., Tinti, C., Spano, P. and Memo, M., Rat pituitary cells selectively express mRNA encoding the short isoform of the γ2 GABA_A receptor subunit, 145

Waeber, C., see Palacios, J.M., 175 Walther, D., see Shimada, S., 359

Wang, C., Li, Y., Wible, B., Angelides, K.J. and Ishii, D.N., Effects of insulin and insulin-like growth factors on neurofilament mRNA and tubulin mRNA content in human neuroblastoma SH-SY5Y cells, 289

Weiss, S.R.B., see Wong, M.-L., 19 Wible, B., see Wang, C., 289

Widdop, R.E., see Rutherfurd, S.D., 301 Williams, C., see Klempt, N.D., 93

Wong, M.-L., Weiss, S.R.B., Gold, P.W., Doi, S.Q., Banerjee, S., Licinio, J., Lad, R., Post, R.M.and Smith, M.A., Induction of constitutive heat shock protein 73 mRNA in the dentate gyrus by seizures, 19

Wooten, G.F., see Pollack, A.E., 35 Worley, P.F., see Bhat, R.V., 263

Yamada, N., see Dragunow, M., 119 Yoong, L.K.K., see Sutherland, M.K., 239 Young III, W.S. and Lightman, S.L., Chronic stress elevates enkephalin expression in the rat paraventricular and supraoptic nuclei, 111

Zoeiler, R.T. and Rudeen, P.K., Ethanol blocks the cold-induced increase in thyrotropin-releasing hormone mRNA in paraventricular nuclei but not the cold-induced increase in thyrotropin, 321

